

EXHIBIT 4

OLSSON, FRANK AND WEEDA, P.C.

PHILIP C. OLSSON
RICHARD L. FRANK
DAVID F. WEEDA (1948-2001)
DENNIS R. JOHNSON
ARTHUR Y. TSIEH
JOHN W. BODE*
STEPHEN D. TERMAN
MARSHALL L. MATZ
MICHAEL J. O'FLAHERTY
DAVID L. DURKIN
NEIL F. O'FLAHERTY
PAMELA J. FURMAN
BRETT T. SCHWEMER
TISH E. PAHL
ROBERT A. HAHN

ATTORNEYS AT LAW
SUITE 400
1400 SIXTEENTH STREET, N.W.
WASHINGTON, D. C. 20036-2220
(202) 789-1212
FACSIMILE (202) 234-3550

STEPHEN L. LACEY
EVAN P. PHELPS
VALERIE B. SOLOMON
JOLYDA O. SWAIM
KATHRYN E. BALMFORD
COUNSEL
NAOMI J. L. HALPERN
OF COUNSEL
JUR. T. STROBOS
JACQUELINE H. EAGLE
KENNETH D. ACKERMAN
MARK L. ITZKOFF
DAVID A. BIEGING
SR. GOVERNMENT AFFAIRS ADVISOR
JOHN R. BLOCK
CHARLES W. STENHOLM
BRIAN E. JOHNSON
SALLY S. DONNER

May 18, 2005

*PRACTICE WITHIN THE DISTRICT OF COLUMBIA
IS LIMITED TO MATTERS AND PROCEEDINGS
BEFORE FEDERAL COURTS AND AGENCIES

Dockets Management Branch (HFA-305)
Food and Drug Administration
Department of Health and Human Services
5630 Fishers Lane
Room 1061
Rockville, MD 20852

Re: Docket No. 2005P-0127 -- ANDAs Referencing Arava® (Leflunomide) Tablets

Dear Food and Drug Administration:

This comment is submitted on behalf of the sponsor of a pending ANDA for 10mg and 20mg generic leflunomide tablets, based on the reference listed drug Arava. This comment responds to the citizen petition submitted by Covington & Burling on behalf of Aventis Pharmaceuticals, Inc., the sponsor of the NDA for Arava, that seeks to raise obstacles to the approval of ANDAs for generic versions of Arava. For the reasons discussed below, the petition is without merit and should be summarily denied.

The crux of Aventis's petition is that FDA should not approve an ANDA for 10mg and 20mg generic leflunomide tablets unless the sponsor demonstrates that five 20mg generic leflunomide tablets are bioequivalent to a single 100mg Arava tablet. Aventis asserts a threefold basis for its request. First, five 20mg Arava tablets have not been shown to be bioequivalent to a single 100mg Arava tablet. Second, the FDA-approved labeling for Arava recommends the use of 100mg Arava tablets as a "loading dose" when an individual is started on Arava therapy. (The maintenance dose of Arava is 20mg daily, or 10mg daily if a 20mg daily dose is not well tolerated clinically.) Third, omission of the "loading dose" labeling information may render a generic product less effective than Arava, thereby precluding ANDA approval.

2005P-0127

C 2

OLSSON, FRANK AND WEEDA, P. C.

Letter to Dockets Management Branch
May 18, 2005
Page 2

On its face, the petition seems to be credible. However, the petition is undercut by the following information in the public domain showing that Aventis discontinued 100mg Arava tablets over three years ago:

- By letter dated January 4, 2002 entitled "Dear Pharmaceutical Buyer," Aventis announced that 100mg Arava tablets had been discontinued. The letter stated: "We will continue to offer Arava in the 10- and 20- mg tablet size." (Exhibit A; available at <http://aventis.custservices.com/news.asp?up=103> (accessed May 16, 2005).)
- By letter dated April 10, 2003, FDA's Office of Generic Drugs notified our client that Aventis had discontinued 100mg Arava tablets. (Exhibit B; client name and portions of letter not relevant to this comment redacted.)
- We are informed that IMS data do not show any sales of 100mg Arava tablets in 2004.

Even though Aventis discontinued 100mg Arava tablets in early 2002, the current professional labeling for Arava (Exhibit C; November 2004 revision, obtained from www.arava.com (accessed May 16, 2005)) nevertheless continues to state that 100mg Arava tablets are available, and that a "loading dose" using 100mg Arava tablets is recommended for patients beginning Arava therapy.

In light of the public information listed above, it should be obvious that the only purpose of the citizen petition submitted on behalf of Aventis is to block, or at least delay, generic competition for Arava. There is no public health basis for the petition.

Taking Aventis's petition at face value, only 100mg Arava tablets – and not five 20mg generic leflunomide tablets or even five 20mg Arava tablets – are appropriate for the "loading doses." Aventis asserts this because five 20mg Arava tablets have not been shown to be bioequivalent to one 100mg Arava tablet. Even though Aventis no longer manufactures and markets 100mg Arava tablets, Aventis's labeling continues to recommend the use of 100mg Arava tablets for the 100mg "loading dose" – a dose that can be reached only by using a combination of 10mg and/or 20mg Arava tablets. However, if 10mg and 20mg Arava tablets are safe and effective for use under the conditions set forth in the current Arava labeling – including use for the "loading dose" – it

OLSSON, FRANK AND WEEDA, P. C.

Letter to Dockets Management Branch
May 18, 2005
Page 3

follows that 10mg and 20mg generic leflunomide tablets otherwise eligible for approval would also be safe and effective under the same conditions of use.¹


In an effort to clothe its petition with an aura of scientific and medical legitimacy, Aventis cites a number of references to support the desirability of rapid attainment of steady state plasma levels of leflunomide and the importance of early therapeutic intervention for rheumatoid arthritis. These purported medical and scientific arguments are nothing more than a smoke screen that should not be allowed to mask the frivolous nature of Aventis's petition. In fact, taking Aventis's arguments at face value, lower-cost, generic 10mg and 20mg leflunomide tablets that have been shown to be bioequivalent to 10mg and 20mg Arava tablets will help American physicians, pharmacists, and patients achieve these desirable goals.

For these reasons, it should be evident that the petition submitted on behalf of Aventis is without merit. Acting Commissioner Lester M. Crawford has recently stated that the Agency will try to improve its decision-making timeframe for citizen petitions, especially those that seek to block or delay generic drug approvals. Consistent with Dr. Crawford's stated intentions, the Agency should summarily deny this citizen petition at once.

Finally, the public information summarized above – none of which was disclosed in the citizen petition – is clearly unfavorable to the petition. This information was clearly known to Aventis, and was presumably known to the signers of the petition, two lawyers from a well-recognized law firm with long-standing expertise in drug regulatory issues. Such a situation is at odds with the petition's certification – pursuant to 21 C.F.R. § 10.30 – that, to the best knowledge and belief of the petitioner, the petition includes information known to the petitioner “which are unfavorable to the petition.”

We appreciate this opportunity to comment.

Respectfully submitted,



Arthur Y. Tsien

Exhibits

¹ The flip side of this coin is that if proposed 10mg and 20mg generic leflunomide tablets cannot be used safely and effectively as the initial “loading dose” under the currently approved labeling for Arava, it follows that 10mg and 20mg Arava tablets also cannot be used safely and effectively for the same purpose under that labeling. Thus, Arava should be withdrawn from the marketplace.

EXHIBIT A

News

Update

Aventis Pharmaceuticals - Arava® 100mg Tablets Discontinued



January 4, 2002

Dear Pharmaceutical Buyer:

Aventis Pharmaceuticals has made a decision to discontinue 100 mg Arava® (leflunomide) tablets trade package. Following is information on the affected product:

Product Description/Size	NDC
100 mg, 3-ct blister pack	0088-2162-03

This discontinuation will take effect immediately and will affect only this trade size. We will continue to offer Arava in the 10- and 20- mg tablet size.

For returns on this product, note that if the product dating falls within the expiration date of the Aventis return policy, then return product to our designated return service, One-Box. If the product dating falls outside of the Aventis return policy dating, then contact our Customer Service Department for return authorization at 1-800-207-8049.

If you have any questions please feel free to contact your Aventis Pharmaceuticals Senior National Trade Account Manager or our Customer Service Department.

Sincerely,

Guerdon R. Green
Director, Trade Administration & Development
Managed Healthcare

Updated :01/04/2002

Current Updates

EXHIBIT B



DEPARTMENT OF HEALTH & HUMAN SERVICES

APR 14 2003

Food and Drug Administration
Rockville MD 20857

APR 10 2003

Reference Number: OGD #02-008

Dear Mr.

This letter is in response to your correspondence dated January 7, 2002. You request that the Office of Generic Drugs (OGD) provide bioequivalence recommendations regarding Leflunomide Tablets, 20 mg. OGD provides the following comments:

1. The designated reference listed drug (RLD) for leflunomide is the 20 mg strength tablets of Arava®. The 100 mg strength has been discontinued from the market. If you wish to submit an application for leflunomide tablets, 100 mg, please submit a Citizen Petition to the Food and Drug Administration (FDA) pursuant to 21 C.F.R. §314.122 to seek a determination whether the said tablets were discontinued for reasons of safety or effectiveness. Please follow the citizen petition format as outlined in 21 C.F.R. §§10.20 and 10.30. If the FDA determines that they were not withdrawn for safety or effectiveness reasons, an additional study on the 100 mg strength will be necessary.

2.

3.


4.

5.

6.

If you have any questions, please call Nina Nwaba, Pharm.D., Project Manager, Division of Bioequivalence, at (301) 827-5847. In future correspondence regarding this issue, please include a copy of this letter.

Sincerely yours,



Gary J. Buehler
Director
Office of Generic Drugs
Center for Drug Evaluation and Research

EXHIBIT C

Rev. November 2004

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg


Rx only

CONTRAINDICATIONS AND WARNINGS

PREGNANCY MUST BE EXCLUDED BEFORE THE START OF TREATMENT WITH ARAVA. ARAVA IS CONTRAINDICATED IN PREGNANT WOMEN, OR WOMEN OF CHILDBEARING POTENTIAL WHO ARE NOT USING RELIABLE CONTRACEPTION. (SEE CONTRAINDICATIONS AND WARNINGS.) PREGNANCY MUST BE AVOIDED DURING ARAVA TREATMENT OR PRIOR TO THE COMPLETION OF THE DRUG ELIMINATION PROCEDURE AFTER ARAVA TREATMENT.

DESCRIPTION

ARAVA® (leflunomide) is a pyrimidine synthesis inhibitor. The chemical name for leflunomide is N-(4'-trifluoromethylphenyl)-5-methylisoxazole-4-carboxamide. It has an empirical formula $C_{12}H_9F_3N_2O_2$, a molecular weight of 270.2 and the following structural formula:




ARAVA is available for oral administration as tablets containing 10, 20, or 100 mg of active drug. Combined with leflunomide are the following inactive ingredients: colloidal silicon dioxide, crospovidone, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, starch, talc, titanium dioxide, and yellow ferric oxide (20 mg tablet only).

CLINICAL PHARMACOLOGY**Mechanism of Action**

Leflunomide is an isoxazole immunomodulatory agent which inhibits dihydroorotate dehydrogenase (an enzyme involved in de novo pyrimidine synthesis) and has antiproliferative activity. Several *in vivo* and *in vitro* experimental models have demonstrated an anti-inflammatory effect.

Pharmacokinetics

Following oral administration, leflunomide is metabolized to an active metabolite A77 1726 (hereafter referred to as M1) which is responsible for essentially all of its activity *in vivo*. Plasma levels of leflunomide are occasionally seen, at very low levels. Studies of the pharmacokinetics of leflunomide have primarily examined the plasma concentrations of this active metabolite.



A77 1726 (M1)

Absorption

Following oral administration, peak levels of the active metabolite, M1, occurred between 6 - 12 hours after dosing. Due to the very long half-life of M1 (~2 weeks), a loading dose of 100 mg for 3 days was used in clinical studies to facilitate the rapid attainment of steady-state levels of M1. Without a loading dose, it is estimated that attainment of steady-state plasma concentrations would require nearly two months of dosing. The resulting plasma concentrations following both loading doses and continued clinical dosing indicate that M1 plasma levels are dose proportional.

Table 1. Pharmacokinetic Parameters for M1 after Administration of Leflunomide at Doses of 5, 10, and 25 mg/day for 24 Weeks to Patients (n=54) with Rheumatoid Arthritis (Mean ± SD) (Study YU204)

Maintenance (Loading) Dose			
Parameter	5 mg (50 mg)	10 mg (100 mg)	25 mg (100 mg)
C_{24} (Day 1) ($\mu\text{g/mL}$) ¹	4.0 ± 0.6	8.4 ± 2.1	8.5 ± 2.2
C_{24} (ss) ($\mu\text{g/mL}$) ²	8.8 ± 2.9	18 ± 9.6	63 ± 36
$t_{1/2}$ (DAYS)	15 ± 3	14 ± 5	18 ± 9

¹ Concentration at 24 hours after loading dose

² Concentration at 24 hours after maintenance doses at steady state

Relative to an oral solution, ARAVA tablets are 80% bioavailable. Co-administration of leflunomide tablets with a high fat meal did not have a significant impact on M1 plasma levels.

Distribution

M1 has a low volume of distribution ($V_{ss} = 0.13$ L/kg) and is extensively bound (>99.3%) to albumin in healthy subjects. Protein binding has been shown to be linear at therapeutic concentrations. The free fraction of M1 is slightly higher in patients with rheumatoid arthritis and approximately doubled in patients with chronic renal failure; the mechanism and significance of these increases are unknown.

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

Metabolism

Leflunomide is metabolized to one primary (M1) and many minor metabolites. Of these minor metabolites, only 4-trifluoromethylaniline (TFMA) is quantifiable, occurring at low levels in the plasma of some patients. The parent compound is rarely detectable in plasma. At the present time the specific site of leflunomide metabolism is unknown. *In vivo* and *in vitro* studies suggest a role for both the GI wall and the liver in drug metabolism. No specific enzyme has been identified as the primary route of metabolism for leflunomide; however, hepatic cytosolic and microsomal cellular fractions have been identified as sites of drug metabolism.

Elimination

The active metabolite M1 is eliminated by further metabolism and subsequent renal excretion as well as by direct biliary excretion. In a 28 day study of drug elimination (n=3) using a single dose of radiolabeled compound, approximately 43% of the total radioactivity was eliminated in the urine and 48% was eliminated in the feces. Subsequent analysis of the samples revealed the primary urinary metabolites to be leflunomide glucuronides and an oxanilic acid derivative of M1. The primary fecal metabolite was M1. Of these two routes of elimination, renal elimination is more significant over the first 96 hours after which fecal elimination begins to predominate. In a study involving the intravenous administration of M1, the clearance was estimated to be 31 mL/hr.

In small studies using activated charcoal (n=1) or cholestyramine (n=3) to facilitate drug elimination, the *in vivo* plasma half-life of M1 was reduced from >1 week to approximately 1 day. (See **PRECAUTIONS – General – Need for Drug Elimination**). Similar reductions in plasma half-life were observed for a series of volunteers (n=96) enrolled in pharmacokinetic trials who were given cholestyramine. This suggests that biliary recycling is a major contributor to the long elimination half-life of M1. Studies with both hemodialysis and CAPD (chronic ambulatory peritoneal dialysis) indicate that M1 is not dialyzable.

Special Populations

Gender. Gender has not been shown to cause a consistent change in the *in vivo* pharmacokinetics of M1.

Age. Age has been shown to cause a change in the *in vivo* pharmacokinetics of M1 (see **CLINICAL PHARMACOLOGY – Special Populations – Pediatrics**).

Smoking. A population based pharmacokinetic analysis of the phase III data indicates that smokers have a 38% increase in clearance over non-smokers; however, no difference in clinical efficacy was seen between smokers and nonsmokers.

Chronic Renal Insufficiency. In single dose studies in patients (n=6) with chronic renal insufficiency requiring either chronic ambulatory peritoneal dialysis (CAPD) or hemodialysis, neither had a significant impact on circulating levels of M1. The free fraction of M1 was almost doubled, but the mechanism of this increase is not known. In light of the fact that the kidney plays a role in drug elimination, and without adequate studies of leflunomide use in subjects with renal insufficiency, caution should be used when ARAVA is administered to these patients.

Hepatic Insufficiency. Studies of the effect of hepatic insufficiency on M1 pharmacokinetics have not been done. Given the need to metabolize leflunomide into the active species, the role of the liver in drug elimination/recycling, and the possible risk of increased hepatic toxicity, the use of leflunomide in patients with hepatic insufficiency is not recommended.

Pediatrics

The pharmacokinetics of M1 following oral administration of leflunomide have been investigated in 73 pediatric patients with polyarticular course Juvenile Rheumatoid Arthritis (JRA) who ranged in age from 3 to 17 years. The results of a population pharmacokinetic analysis of these trials have demonstrated that pediatric patients with body weights ≤ 40 kg have a reduced clearance of M1 (see Table 2) relative to adult rheumatoid arthritis patients.

Table 2: Population Pharmacokinetic Estimate of M1 Clearance Following Oral Administration of Leflunomide in Pediatric Patients with Polyarticular Course JRA Mean \pm SD [Range]

N	Body Weight (kg)	CL (mL/h)
10	<20	18 \pm 9.8 [6.8-37]
30	20-40	18 \pm 9.5 [4.2-43]
33	>40	26 \pm 16 [9.7-93.6]

Drug Interactions

In vivo drug interaction studies have demonstrated a lack of a significant drug interaction between leflunomide and tri-phasic oral contraceptives, and cimetidine.

In vitro studies of protein binding indicated that warfarin did not affect M1 protein binding. At the same time M1 was shown to cause increases ranging from 13 - 50% in the free fraction of diclofenac, ibuprofen and tolbutamide at concentrations in the clinical range. *In vitro* studies of drug metabolism indicate that M1 inhibits CYP 450 2C9, which is responsible for the metabolism of phenytoin, tolbutamide, warfarin and many NSAIDs. M1 has been shown to inhibit the formation of 4-hydroxydiclofenac from diclofenac *in vitro*. The clinical significance of these findings with regard to phenytoin and tolbutamide is unknown, however, there was extensive concomitant use of NSAIDs in the clinical studies and no differential effect was observed. (See **PRECAUTIONS – Drug Interactions**).

Methotrexate. Coadministration, in 30 patients, of ARAVA (100 mg/day x 2 days followed by 10 - 20 mg/day) with methotrexate (10 - 25 mg/week, with folate) demonstrated no pharmacokinetic interaction between the two drugs. However, co-administration increased risk of hepatotoxicity (see **PRECAUTIONS – Drug Interactions – Hepatotoxic Drugs**).

Rifampin. Following concomitant administration of a single dose of ARAVA to subjects receiving multiple doses of rifampin, M1 peak levels were increased (~40%) over those seen when ARAVA was given alone. Because of the potential for ARAVA levels to continue to increase with multiple dosing, caution should be used if patients are to receive both ARAVA and rifampin.

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

CLINICAL STUDIES

A. ADULTS

The efficacy of ARAVA in the treatment of rheumatoid arthritis (RA) was demonstrated in three controlled trials showing reduction in signs and symptoms, and inhibition of structural damage. In two placebo controlled trials, efficacy was demonstrated for improvement in physical function.

1. Reduction of signs and symptoms

Relief of signs and symptoms was assessed using the American College of Rheumatology (ACR)20 Responder Index, a composite of clinical, laboratory, and functional measures in rheumatoid arthritis. An "ACR20 Responder" is a patient who had $\geq 20\%$ improvement in both tender and swollen joint counts and in 3 of the following 5 criteria: physician global assessment, patient global assessment, functional ability measure [Modified Health Assessment Questionnaire (MHAQ)], visual analog pain scale, and erythrocyte sedimentation rate or C-reactive protein. An "ACR20 Responder at Endpoint" is a patient who completed the study and was an ACR20 Responder at the completion of the study.

2. Inhibition of structural damage

Inhibition of structural damage compared to control was assessed using the Sharp Score (Sharp, JT. Scoring Radiographic Abnormalities in Rheumatoid Arthritis, Radiologic Clinics of North America, 1996; vol. 34, pp. 233-241), a composite score of X-ray erosions and joint space narrowing in hands/wrists and forefeet.

3. Improvement in physical function

Improvement in physical function was assessed using the Health Assessment Questionnaire (HAQ) and the Medical Outcomes Survey Short Form (SF-36). In all Arava monotherapy studies, an initial loading dose of 100 mg per day for three days only was used followed by 20 mg per day thereafter.

US301 Clinical Trial in Adults

Study US301, a 2 year study, randomized 482 patients with active RA of at least 6 months duration to leflunomide 20 mg/day (n=182), methotrexate 7.5 mg/week increasing to 15 mg/week (n=182), or placebo (n=118). All patients received folate 1 mg BID. Primary analysis was at 52 weeks with blinded treatment to 104 weeks.

Overall, 235 of the 508 randomized treated patients (482 in primary data analysis and an additional 26 patients), continued into a second 12 months of double-blind treatment (98 leflunomide, 101 methotrexate, 36 placebo). Leflunomide dose continued at 20 mg/day and the methotrexate dose could be increased to a maximum of 20 mg/week. In total, 190 patients (83 leflunomide, 80 methotrexate, 27 placebo) completed 2 years of double-blind treatment.

The rate and reason for withdrawal is summarized in Table 3.

Table 3: Withdrawals in US301

	n(%) patients		
	Leflunomide 190	Placebo 128	Methotrexate 190
Withdrawals in Year-1			
Lack of efficacy	33 (17.4)	70 (54.7)	50 (26.3)
Safety	44 (23.2)	12 (9.4)	22 (11.6)
Other ¹	15 (7.9)	10 (7.8)	17 (9.0)
Total	92 (48.4)	92 (71.9)	89 (46.8)
Patients entering Year 2	98	36	101
Withdrawals in Year-2			
Lack of efficacy	4 (4.1)	1 (2.8)	4 (4.0)
Safety	8 (8.2)	0 (0.0)	10 (9.9)
Other ¹	3 (3.1)	8 (22.2)	7 (6.9)
Total	15 (15.3)	9 (25.0)	21 (20.8)

¹ Includes: lost to follow up, protocol violation, noncompliance, voluntary withdrawal, investigator discretion.

MN301/303/305 Clinical Trial in Adults

Study MN301 randomized 358 patients with active RA to leflunomide 20 mg/day (n=133), sulfasalazine 2.0 g/day (n=133), or placebo (n=92). Treatment duration was 24 weeks. An extension of the study was an optional 6-month blinded continuation of MN301 without the placebo arm, resulting in a 12-month comparison of leflunomide and sulfasalazine (study MN303).

Of the 168 patients who completed 12 months of treatment in MN301 and MN303, 146 patients (87%) entered a 1-year extension study of double blind active treatment (MN305; 60 leflunomide, 60 sulfasalazine, 26 placebo/ sulfasalazine). Patients continued on the same daily dosage of leflunomide or sulfasalazine that they had been taking at the completion of MN301/303. A total of 121 patients (53 leflunomide, 47 sulfasalazine, 21 placebo/ sulfasalazine) completed the 2 years of double-blind treatment.

Patient withdrawal data in MN301/303/305 is summarized in Table 4.

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

Table 4: Withdrawals in study MN301/303/305

	n(%) patients		
	Leflunomide 133	Placebo 92	Sulfasalazine 133
Withdrawals in MN301 (Mo 0-6)			
Lack of efficacy	10 (7.5)	29 (31.5)	14 (10.5)
Safety	19 (14.3)	6 (6.5)	25 (18.8)
Other ¹	8 (6.0)	6 (6.5)	11 (8.3)
Total	37 (27.8)	41 (44.6)	50 (37.6)
Patients entering MN303	80		76
Withdrawals in MN303 (Mo 7-12)			
Lack of efficacy	4 (5.0)		2 (2.6)
Safety	2 (2.5)		5 (6.6)
Other ¹	3 (3.8)		1 (1.3)
Total	9 (11.3)		8 (10.5)
Patients entering MN305	60		60
Withdrawals in MN305 (Mo 13-24)			
Lack of efficacy	0 (0.0)		3 (5.0)
Safety	6 (10.0)		8 (13.3)
Other ¹	1 (1.7)		2 (3.3)
Total	7 (11.7)		13 (21.7)

¹ Includes: lost to follow up, protocol violation, noncompliance, voluntary withdrawal, investigator discretion.

MN302/304 Clinical Trial in Adults

Study MN302 randomized 999 patients with active RA to leflunomide 20 mg/day (n=501) or methotrexate at 7.5 mg/week increasing to 15 mg/week (n=498). Folate supplementation was used in 10% of patients. Treatment duration was 52 weeks.

Of the 736 patients who completed 52 weeks of treatment in study MN302, 612 (83%) entered the double-blind, 1-year extension study MN304 (292 leflunomide, 320 methotrexate). Patients continued on the same daily dosage of leflunomide or methotrexate that they had been taking at the completion of MN302. There were 533 patients (256 leflunomide, 277 methotrexate) who completed 2 years of double-blind treatment.

Patient withdrawal data in MN302/304 is summarized in Table 5.

Table 5: Withdrawals in MN302/304

	n(%) patients	
	Leflunomide 501	Methotrexate 498
Withdrawals in MN302 (Year-1)		
Lack of efficacy	37 (7.4)	15 (3.0)
Safety	98 (19.6)	79 (15.9)
Other ¹	17 (3.4)	17 (3.4)
Total	152 (30.3)	111 (22.3)
Patients entering MN304	292	320
Withdrawals in MN304 (Year-2)		
Lack of efficacy	13 (4.5)	9 (2.8)
Safety	11 (3.8)	22 (6.9)
Other ¹	12 (4.1)	12 (3.8)
Total	36 (12.3)	43 (13.4)


¹ Includes: lost to follow up, protocol violation, noncompliance, voluntary withdrawal, investigator discretion.

Clinical Trial Data**1. Signs and symptoms Rheumatoid Arthritis**

The ACR20 Responder at Endpoint rates are shown in Figure 1. ARAVA was statistically significantly superior to placebo in reducing the signs and symptoms of RA by the primary efficacy analysis, ACR20 Responder at Endpoint, in study US301 (at the primary 12 months endpoint) and MN301 (at 6 month endpoint). ACR20 Responder at Endpoint rates with ARAVA treatment were consistent across the 6 and 12 month studies (41 - 49%). No consistent differences were demonstrated between leflunomide and methotrexate or between leflunomide and sulfasalazine. ARAVA treatment effect was evident by 1 month, stabilized by 3 - 6 months, and continued throughout the course of treatment as shown in Figure 2.


ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

Figure 1



	Comparisons	95% Confidence Interval	p Value
US301	Leflunomide vs. Placebo	(12, 32)	<0.0001
	Methotrexate vs. Placebo	(8, 30)	<0.0001
	Leflunomide vs. Methotrexate	(-4, 16)	NS
MN301	Leflunomide vs. Placebo	(7, 33)	0.0026
	Sulfasalazine vs. Placebo	(4, 29)	0.0121
	Leflunomide vs. Sulfasalazine	(-8, 16)	NS
MN302	Leflunomide vs. Methotrexate	(-19, -7)	<0.0001

Figure 2



*Last Observation Carried Forward.

10 mg, 20 mg, 100 mg

ACR50 and ACR70 Responders are defined in an analogous manner to the ACR 20 Responder, but use improvements of 50% or 70%, respectively (Table 6). Mean change for the individual components of the ACR Responder Index are shown in Table 7.

Table 6. Summary of ACR Response Rates*

Study and Treatment Group	ACR20	ACR50	ACR70
Placebo-Controlled Studies			
US301 (12 months)			
Leflunomide (n=178) [†]	52.2 [‡]	34.3 [‡]	20.2 [‡]
Placebo (n=118) [†]	26.3	7.6	4.2
Methotrexate (n=180) [†]	45.6	22.8	9.4
MN301 (6 months)			
Leflunomide (n=130) [†]	54.6 [‡]	33.1 [‡]	10.0 [§]
Placebo (n=91) [†]	28.6	14.3	2.2
Sulfasalazine (n=132) [†]	56.8	30.3	7.6
Non-Placebo Active-Controlled Studies			
MN302 (12 months)			
Leflunomide (n=495) [†]	51.1	31.1	9.9
Methotrexate (n=489) [†]	65.2	43.8	16.4

* Intent to treat (ITT) analysis using last observation carried forward (LOCF) technique for patients who discontinued early.
[†] N is the number of ITT patients for whom adequate data were available to calculate the indicated rates.
[‡] p<0.001 leflunomide vs placebo
[§] p<0.02 leflunomide vs placebo

Table 7 shows the results of the components of the ACR response criteria for US301, MN301, and MN302. ARAVA was significantly superior to placebo in all components of the ACR Response criteria in study US301 and MN301. In addition, Arava was significantly superior to placebo in improving morning stiffness, a measure of RA disease activity, not included in the ACR Response criteria. No consistent differences were demonstrated between ARAVA and the active comparators.

Table 7. Mean Change in the Components of the ACR Responder Index*

Components	Placebo-Controlled Studies						Non-placebo Controlled Study	
	US301 (12 months)			MN301 Non-US (6 months)			MN302 Non-US (12 months)	
	Leflunomide	Methotrexate	Placebo	Leflunomide	Sulfasalazine	Placebo	Leflunomide	Methotrexate
Tender joint count ¹	-7.7	-6.6	-3.0	-9.7	-8.1	-4.3	-8.3	-9.7
Swollen joint count ¹	-5.7	-5.4	-2.9	-7.2	-6.2	-3.4	-6.8	-9.0
Patient global assessment ²	-2.1	-1.5	0.1	-2.8	-2.6	-0.9	-2.3	-3.0
Physician global assessment ²	-2.8	-2.4	-1.0	-2.7	-2.5	-0.8	-2.3	-3.1
Physical function/disability (MHAQ/HAQ)	-0.29	-0.15	0.07	-0.50	-0.29	-0.04	-0.37	-0.44
Pain intensity ²	-2.2	-1.7	-0.5	-2.7	-2.0	-0.9	-2.1	-2.9
Erythrocyte Sedimentation rate	-6.26	-6.48	2.56	-7.48	-16.56	3.44	-10.12	-22.18
C-reactive protein	-0.62	-0.50	0.47	-2.26	-1.19	0.16	-1.86	-2.45
Not included in the ACR Responder Index								
Morning Stiffness (min)	-101.4	-88.7	14.7	-93.0	-42.4	-6.8	-63.7	-86.6

* Last Observation Carried Forward; Negative Change Indicates Improvement

¹ Based on 28 joint count

² Visual Analog Scale - 0=Best; 10=Worst

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

Maintenance of effect


After completing 12 months of treatment, patients continuing on study treatment were evaluated for an additional 12 months of double-blind treatment (total treatment period of 2 years) in studies US301, MN305, and MN304. ACR Responder rates at 12 months were maintained over 2 years in most patients continuing a second year of treatment.

Improvement from baseline in the individual components of the ACR responder criteria was also sustained in most patients during the second year of Arava treatment in all three trials.

2. Inhibition of structural damage

The change from baseline to endpoint in progression of structural disease, as measured by the Sharp X-ray score, is displayed in Figure 3. ARAVA was statistically significantly superior to placebo in inhibiting the progression of disease by the Sharp Score. No consistent differences were demonstrated between leflunomide and methotrexate or between leflunomide and sulfasalazine.

Figure 3



L= Leflunomide; M=methotrexate; S=sulfasalazine; P=placebo

	Comparisons	95% Confidence Interval	p Value
US301	Leflunomide vs. Placebo	(-4.0, -1.1)	0.0007
	Methotrexate vs. Placebo	(-2.6, -0.2)	0.0196
	Leflunomide vs. Methotrexate	(-2.3, 0.0)	0.0499
MN301	Leflunomide vs. Placebo	(-6.2, -1.8)	0.0004
	Sulfasalazine vs. Placebo	(-6.9, 0.0)	0.0484
	Leflunomide vs. Sulfasalazine	(-3.3, 1.2)	NS
MN302	Leflunomide vs. Methotrexate	(-2.2, 7.4)	NS


3. Improvement in physical function

The Health Assessment Questionnaire (HAQ) assesses a patient's physical function and degree of disability. The mean change from baseline in functional ability as measured by the HAQ Disability Index (HAQ DI) in the 6 and 12 month placebo and active controlled trials is shown in Figure 4. ARAVA was statistically significantly superior to placebo in improving physical function. Superiority to placebo was demonstrated consistently across all eight HAQ DI subscales (dressing, arising, eating, walking, hygiene, reach, grip and activities) in both placebo controlled studies.

The Medical Outcomes Survey Short Form 36 (SF-36), a generic health-related quality of life questionnaire, further addresses physical function. In US301, at 12 months, ARAVA provided statistically significant improvements compared to placebo in the Physical Component Summary (PCS) Score.

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

Figure 4



* as measured by HAQ Disability Index

L= Leflunomide, M=Methotrexate, P=Placebo, S=Sulfasalazine

	Comparison	95% Confidence Interval	p Value
US301	Leflunomide vs. Placebo	(-0.58, -0.29)	0.0001
	Leflunomide vs. Methotrexate	(-0.34, -0.07)	0.0026
MN301	Leflunomide vs. Placebo	(-0.67, -0.36)	<0.0001
	Leflunomide vs. Sulfasalazine	(-0.33, -0.03)	0.0163
MN302	Leflunomide vs. Methotrexate	(0.01, 0.16)	0.0221

Maintenance of effect

The improvement in physical function demonstrated at 6 and 12 months was maintained over two years. In those patients continuing therapy for a second year, this improvement in physical function as measured by HAQ and SF-36 (PCS) was maintained.

B. PEDIATRICS**Clinical Trials in Pediatrics**

ARAVA was studied in a single multicenter, double-blind, active-controlled trial in 94 patients (1:1 randomization) with polyarticular course juvenile rheumatoid arthritis (JRA) as defined by the American College of Rheumatology (ACR). Approximately 68% of pediatric patients receiving ARAVA, versus 89% of pediatric patients receiving the active comparator, improved by Week 16 (end-of-study) employing the JRA Definition of Improvement (DOI) $\geq 30\%$ responder endpoint. In this trial, the loading dose and maintenance dose of ARAVA was based on three weight categories: <20 kg, 20-40kg, and ≥ 40 kg. The response rate to ARAVA in pediatric patients ≤ 40 kg was less robust than in pediatric patients >40 kg suggesting suboptimal dosing in smaller weight pediatric patients, as studied, resulting in less than efficacious plasma concentrations, despite reduced clearance of M1. (See **PHARMACOKINETICS – Pediatrics**).

INDICATIONS AND USAGE

ARAVA is indicated in adults for the treatment of active rheumatoid arthritis (RA):

1. to reduce signs and symptoms
2. to inhibit structural damage as evidenced by X-ray erosions and joint space narrowing
3. to improve physical function.

(see **CLINICAL STUDIES**)

Aspirin, nonsteroidal anti-inflammatory agents and/or low dose corticosteroids may be continued during treatment with ARAVA (see **PRECAUTIONS – Drug Interactions – NSAIDs**). The combined use of ARAVA with antimalarials, intramuscular or oral gold, D penicillamine, azathioprine, or methotrexate has not been adequately studied. (See **WARNINGS – Immunosuppression Potential/Bone Marrow Suppression**).

CONTRAINDICATIONS

ARAVA is contraindicated in patients with known hypersensitivity to leflunomide or any of the other components of ARAVA.

ARAVA can cause fetal harm when administered to a pregnant woman. Leflunomide, when administered orally to rats during organogenesis at a dose of 15 mg/kg, was teratogenic (most notably anophthalmia or microphthalmia and internal hydrocephalus). The systemic exposure of rats at this dose was approximately 1/10 the human exposure level based on AUC. Under these exposure conditions, leflunomide also caused a decrease in the maternal body weight and an increase in embryoletality with a decrease in fetal body weight for surviving fetuses. In rabbits, oral treatment with 10 mg/kg of leflunomide during organogenesis resulted in fused, dysplastic sternebrae. The exposure level at this dose was essentially equivalent to the maximum human exposure level based on AUC. At a 1 mg/kg dose, leflunomide was not teratogenic in rats and rabbits.

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

When female rats were treated with 1.25 mg/kg of leflunomide beginning 14 days before mating and continuing until the end of lactation, the offspring exhibited marked (greater than 90%) decreases in postnatal survival. The systemic exposure level at 1.25 mg/kg was approximately 1/100 the human exposure level based on AUC.

ARAVA is contraindicated in women who are or may become pregnant. If this drug is used during pregnancy, or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the fetus.

WARNINGS

Immunosuppression Potential/Bone Marrow Suppression

ARAVA is not recommended for patients with severe immunodeficiency, bone marrow dysplasia, or severe, uncontrolled infections. In the event that a serious infection occurs, it may be necessary to interrupt therapy with ARAVA and administer cholestyramine or charcoal (see **PRECAUTIONS – General – Need for Drug Elimination**). Medications like leflunomide that have immunosuppression potential may cause patients to be more susceptible to infections, including opportunistic infections. Rarely, severe infections including sepsis, which may be fatal, have been reported in patients receiving ARAVA. Most of the reports were confounded by concomitant immunosuppressant therapy and/or comorbid illness which, in addition to rheumatoid disease, may predispose patients to infection.

There have been rare reports of pancytopenia, agranulocytosis and thrombocytopenia in patients receiving ARAVA alone. These events have been reported most frequently in patients who received concomitant treatment with methotrexate or other immunosuppressive agents, or who had recently discontinued these therapies; in some cases, patients had a prior history of a significant hematologic abnormality.

Patients taking ARAVA should have platelet, white blood cell count and hemoglobin or hematocrit monitored at baseline and monthly for six months following initiation of therapy and every 6 to 8 weeks thereafter. If used with concomitant methotrexate and/or other potential immunosuppressive agents, chronic monitoring should be monthly. If evidence of bone marrow suppression occurs in a patient taking ARAVA, treatment with ARAVA should be stopped, and cholestyramine or charcoal should be used to reduce the plasma concentration of leflunomide active metabolite (see **PRECAUTIONS – General – Need for Drug Elimination**).

In any situation in which the decision is made to switch from ARAVA to another anti-rheumatic agent with a known potential for hematologic suppression, it would be prudent to monitor for hematologic toxicity, because there will be overlap of systemic exposure to both compounds. ARAVA washout with cholestyramine or charcoal may decrease this risk, but also may induce disease worsening if the patient had been responding to ARAVA treatment.

Hepatotoxicity

RARE CASES OF SEVERE LIVER INJURY, INCLUDING CASES WITH FATAL OUTCOME, HAVE BEEN REPORTED DURING TREATMENT WITH LEFLUNOMIDE. MOST CASES OF SEVERE LIVER INJURY OCCUR WITHIN 6 MONTHS OF THERAPY AND IN A SETTING OF MULTIPLE RISK FACTORS FOR HEPATOTOXICITY (liver disease, other hepatotoxins). (See PRECAUTIONS).

At minimum, ALT (SGPT) must be performed at baseline and monitored initially at monthly intervals during the first six months then, if stable, every 6 to 8 weeks thereafter. In addition, if ARAVA and methotrexate are given concomitantly, ACR guidelines for monitoring methotrexate liver toxicity must be followed with ALT, AST, and serum albumin testing monthly.

Guidelines for dose adjustment or discontinuation based on the severity and persistence of ALT elevation are recommended as follows: For confirmed ALT elevations between 2- and 3-fold ULN, dose reduction to 10 mg/day may allow continued administration of ARAVA under close monitoring. If elevations between 2- and 3-fold ULN persist despite dose reduction or if ALT elevations of >3-fold ULN are present, ARAVA should be discontinued and cholestyramine or charcoal should be administered (see **PRECAUTIONS – General – Need for Drug Elimination**) with close monitoring, including retreatment with cholestyramine or charcoal as indicated.

In clinical trials, ARAVA treatment as monotherapy or in combination with methotrexate was associated with elevations of liver enzymes, primarily ALT and AST, in a significant number of patients; these effects were generally reversible. Most transaminase elevations were mild (\leq 2-fold ULN) and usually resolved while continuing treatment. Marked elevations (>3-fold ULN) occurred infrequently and reversed with dose reduction or discontinuation of treatment. Table 8 shows liver enzyme elevations seen with monthly monitoring in clinical trials US301 and MN301. It was notable that the absence of folate use in MN302 was associated with a considerably greater incidence of liver enzyme elevation on methotrexate.

Table 8. Liver Enzyme Elevations >3-fold Upper Limits of Normal (ULN)

	US301			MN301			MN302*	
	LEF	PL	MTX	LEF	PL	SSZ	LEF	MTX
ALT (SGPT) >3-fold ULN (n %)	8 (4.4)	3 (2.5)	5 (2.7)	2 (1.5)	1 (1.1)	2 (1.5)	13 (2.6)	83 (16.7)
Reversed to \leq 2-fold ULN:	8	3	5	2	1	2	12	82
Timing of Elevation								
0-3 Months	6	1	1	2	1	2	7	27
4-6 Months	1	1	3	-	-	-	1	34
7-9 Months	1	1	1	-	-	-	-	16
10-12 Months	-	-	-	-	-	-	5	6

*Only 10% of patients in MN302 received folate. All patients in US301 received folate.

In a 6 month study of 263 patients with persistent active rheumatoid arthritis despite methotrexate therapy, and with normal LFTs, leflunomide was added to a group of 133 patients starting at 10 mg per day and increased to 20 mg as needed. An increase in ALT greater than or equal to three times the ULN was observed in 3.8% of patients compared to 0.8% in 130 patients continued on methotrexate with placebo added.

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

Pre-existing Hepatic Disease

Given the possible risk of increased hepatotoxicity, and the role of the liver in drug activation, elimination and recycling, the use of ARAVA is not recommended in patients with significant hepatic impairment or evidence of infection with hepatitis B or C viruses. (See **Warnings – Hepatotoxicity**).

Skin Reactions

Rare cases of Stevens-Johnson syndrome and toxic epidermal necrolysis have been reported in patients receiving ARAVA. If a patient taking ARAVA develops any of these conditions, ARAVA therapy should be stopped, and a drug elimination procedure is recommended. (See **PRECAUTIONS – General – Need for Drug Elimination**).

Malignancy

The risk of malignancy, particularly lymphoproliferative disorders, is increased with the use of some immunosuppression medications. There is a potential for immunosuppression with ARAVA. No apparent increase in the incidence of malignancies and lymphoproliferative disorders was reported in the clinical trials of ARAVA, but larger and longer-term studies would be needed to determine whether there is an increased risk of malignancy or lymphoproliferative disorders with ARAVA.

Use in Women of Childbearing Potential

There are no adequate and well-controlled studies evaluating ARAVA in pregnant women. However, based on animal studies, leflunomide may increase the risk of fetal death or teratogenic effects when administered to a pregnant woman (see **CONTRAINDICATIONS**). Women of childbearing potential must not be started on ARAVA until pregnancy is excluded and it has been confirmed that they are using reliable contraception. Before starting treatment with ARAVA, patients must be fully counseled on the potential for serious risk to the fetus.

The patient must be advised that if there is any delay in onset of menses or any other reason to suspect pregnancy, they must notify the physician immediately for pregnancy testing and, if positive, the physician and patient must discuss the risk to the pregnancy. It is possible that rapidly lowering the blood level of the active metabolite by instituting the drug elimination procedure described below at the first delay of menses may decrease the risk to the fetus from ARAVA.

Upon discontinuing ARAVA, it is recommended that all women of childbearing potential undergo the drug elimination procedure described below. Women receiving ARAVA treatment who wish to become pregnant must discontinue ARAVA and undergo the drug elimination procedure described below which includes verification of M1 metabolite plasma levels less than 0.02 mg/L (0.02 µg/mL). Human plasma levels of the active metabolite (M1) less than 0.02 mg/L (0.02 µg/mL) are expected to have minimal risk based on available animal data.

Drug Elimination Procedure

The following drug elimination procedure is recommended to achieve non-detectable plasma levels (less than 0.02 mg/L or 0.02 µg/mL) after stopping treatment with ARAVA:

- 1) Administer cholestyramine 8 grams 3 times daily for 11 days. (The 11 days do not need to be consecutive unless there is a need to lower the plasma level rapidly.)
- 2) Verify plasma levels less than 0.02 mg/L (0.02 µg/mL) by two separate tests at least 14 days apart. If plasma levels are higher than 0.02 mg/L, additional cholestyramine treatment should be considered.

Without the drug elimination procedure, it may take up to 2 years to reach plasma M1 metabolite levels less than 0.02 mg/L due to individual variation in drug clearance.

PRECAUTIONS

General

Need for Drug Elimination

The active metabolite of leflunomide is eliminated slowly from the plasma. In instances of any serious toxicity from ARAVA, including hypersensitivity, use of a drug elimination procedure as described in this section is highly recommended to reduce the drug concentration more rapidly after stopping ARAVA therapy. If hypersensitivity is the suspected clinical mechanism, more prolonged cholestyramine or charcoal administration may be necessary to achieve rapid and sufficient clearance. The duration may be modified based on the clinical status of the patient.

Cholestyramine given orally at a dose of 8 g three times a day for 24 hours to three healthy volunteers decreased plasma levels of M1 by approximately 40% in 24 hours and by 49 to 65% in 48 hours.

Administration of activated charcoal (powder made into a suspension) orally or via nasogastric tube (50 g every 6 hours for 24 hours) has been shown to reduce plasma concentrations of the active metabolite, M1, by 37% in 24 hours and by 48% in 48 hours.

These drug elimination procedures may be repeated if clinically necessary.

Respiratory

Interstitial lung disease has been reported during treatment with leflunomide and has been associated with fatal outcomes (see **ADVERSE REACTIONS**). Interstitial lung disease is a potentially fatal disorder, which may occur acutely at any time during therapy and has a variable clinical presentation. New onset or worsening pulmonary symptoms, such as cough and dyspnea, with or without associated fever, may be a reason for discontinuation of the therapy and for further investigation as appropriate. If discontinuation of the drug is necessary, initiation of wash-out procedures should be considered. (See **WARNINGS – Drug Elimination Procedure**).

Renal Insufficiency

Single dose studies in dialysis patients show a doubling of the free fraction of M1 in plasma. There is no clinical experience in the use of ARAVA in patients with renal impairment. Caution should be used when administering this drug in this population.

Vaccinations

No clinical data are available on the efficacy and safety of vaccinations during ARAVA treatment. Vaccination with live vaccines is, however, not recommended. The long half-life of ARAVA should be considered when contemplating administration of a live vaccine after stopping ARAVA.

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

Information for Patients

- The potential for increased risk of birth defects should be discussed with female patients of childbearing potential. It is recommended that physicians advise women that they may be at increased risk of having a child with birth defects if they are pregnant when taking ARAVA, become pregnant while taking ARAVA, or do not wait to become pregnant until they have stopped taking ARAVA and followed the drug elimination procedure (as described in **WARNINGS – Use In Women of Childbearing Potential – Drug Elimination Procedure**).
- Patients should be advised of the possibility of rare, serious skin reactions. Patients should be instructed to inform their physicians promptly if they develop a skin rash or mucous membrane lesions.
- Patients should be advised of the potential hepatotoxic effects of ARAVA and of the need for monitoring liver enzymes. Patients should be instructed to notify their physicians if they develop symptoms such as unusual tiredness, abdominal pain or jaundice.
- Patients should be advised that they may develop a lowering of their blood counts and should have frequent hematologic monitoring. This is particularly important for patients who are receiving other immunosuppressive therapy concurrently with ARAVA, who have recently discontinued such therapy before starting treatment with ARAVA, or who have had a history of a significant hematologic abnormality. Patients should be instructed to notify their physicians promptly if they notice symptoms of pancytopenia (such as easy bruising or bleeding, recurrent infections, fever, paleness or unusual tiredness).
- Patients should be informed about the early warning signs of interstitial lung disease and asked to contact their physician as soon as possible if these symptoms appear or worsen during therapy.

Laboratory Tests

Hematologic Monitoring

At minimum, patients taking ARAVA should have platelet, white blood cell count and hemoglobin or hematocrit monitored at baseline and monthly for six months following initiation of therapy and every 6 to 8 weeks thereafter.

Bone Marrow Suppression Monitoring

If used with concomitantly with immunosuppressants such as methotrexate, chronic monitoring should be monthly. (See **WARNINGS – Immunosuppression Potential/Bone Marrow Suppression**).

Liver Enzyme Monitoring

ALT (SGPT) must be performed at baseline and monitored at monthly intervals during the first six months then, if stable, every 6 to 8 weeks thereafter. In addition, if ARAVA and methotrexate are given concomitantly, ACR guidelines for monitoring methotrexate liver toxicity must be followed with ALT, AST, and serum albumin testing every month. (See **WARNINGS – Hepatotoxicity**.)

Due to a specific effect on the brush border of the renal proximal tubule, ARAVA has a uricosuric effect. A separate effect of hypophosphaturia is seen in some patients. These effects have not been seen together, nor have there been alterations in renal function.

Drug Interactions

Cholestyramine and Charcoal

Administration of cholestyramine or activated charcoal in patients (n=13) and volunteers (n=96) resulted in a rapid and significant decrease in plasma M1 (the active metabolite of leflunomide) concentration (see **PRECAUTIONS – General – Need for Drug Elimination**).

Hepatotoxic Drugs

Increased side effects may occur when leflunomide is given concomitantly with hepatotoxic substances. This is also to be considered when leflunomide treatment is followed by such drugs without a drug elimination procedure. In a small (n=30) combination study of ARAVA with methotrexate, a 2- to 3-fold elevation in liver enzymes was seen in 5 of 30 patients. All elevations resolved, 2 with continuation of both drugs and 3 after discontinuation of leflunomide. A >3-fold increase was seen in another 5 patients. All of these also resolved, 2 with continuation of both drugs and 3 after discontinuation of leflunomide. Three patients met "ACR criteria" for liver biopsy (1: Roenik Grade I, 2: Roenik Grade IIIa). No pharmacokinetic interaction was identified (see **CLINICAL PHARMACOLOGY**).

NSAIDs

In *in vitro* studies, M1 was shown to cause increases ranging from 13 - 50% in the free fraction of diclofenac and ibuprofen at concentrations in the clinical range. The clinical significance of this finding is unknown; however, there was extensive concomitant use of NSAIDs in clinical studies and no differential effect was observed.

Tolbutamide

In *in vitro* studies, M1 was shown to cause increases ranging from 13 - 50% in the free fraction of tolbutamide at concentrations in the clinical range. The clinical significance of this finding is unknown.

Rifampin

Following concomitant administration of a single dose of ARAVA to subjects receiving multiple doses of rifampin, M1 peak levels were increased (~40%) over those seen when ARAVA was given alone. Because of the potential for ARAVA levels to continue to increase with multiple dosing, caution should be used if patients are to be receiving both ARAVA and rifampin.

Warfarin

Increased INR (International Normalized Ratio) when ARAVA and warfarin were co-administered has been rarely reported.

Carcinogenesis, Mutagenesis, and Impairment of Fertility

No evidence of carcinogenicity was observed in a 2-year bioassay in rats at oral doses of leflunomide up to the maximally tolerated dose of 6 mg/kg (approximately 1/40 the maximum human M1 systemic exposure based on AUC). However, male mice in a 2-year bioassay exhibited an increased incidence in lymphoma at an oral dose of 15 mg/kg, the highest dose studied (1.7 times the human M1 exposure based on AUC). Female mice, in the same study, exhibited a dose-related increased incidence of bronchoalveolar adenomas and carcinomas combined beginning at 1.5 mg/kg (approximately 1/10 the human M1 exposure based on AUC). The significance of the findings in mice relative to the clinical use of ARAVA is not known.

Leflunomide was not mutagenic in the Ames Assay, the Unscheduled DNA Synthesis Assay, or in the HGPRT Gene Mutation Assay. In addition, leflunomide was not clastogenic in the *in vivo* Mouse Micronucleus Assay nor in the *in vivo* Cytogenetic Test in Chinese Hamster Bone Marrow Cells. However,

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

4-trifluoromethylaniline (TFMA), a minor metabolite of leflunomide, was mutagenic in the Ames Assay and in the HGPRT Gene Mutation Assay, and was clastogenic in the *in vitro* Assay for Chromosome Aberrations in the Chinese Hamster Cells. TFMA was not clastogenic in the *in vivo* Mouse Micronucleus Assay nor in the *in vivo* Cytogenetic Test in Chinese Hamster Bone Marrow Cells. Leflunomide had no effect on fertility in either male or female rats at oral doses up to 4.0 mg/kg (approximately 1/30 the human M1 exposure based on AUC).

Pregnancy

Pregnancy Category X. (See **CONTRAINDICATIONS** section). Pregnancy Registry: To monitor fetal outcomes of pregnant women exposed to leflunomide, health care providers are encouraged to register such patients by calling 1-877-311-8972.

Nursing Mothers

ARAVA should not be used by nursing mothers. It is not known whether ARAVA is excreted in human milk. Many drugs are excreted in human milk, and there is a potential for serious adverse reactions in nursing infants from ARAVA. Therefore, a decision should be made whether to proceed with nursing or to initiate treatment with ARAVA, taking into account the importance of the drug to the mother.

Use in Males

Available information does not suggest that ARAVA would be associated with an increased risk of male-mediated fetal toxicity. However, animal studies to evaluate this specific risk have not been conducted. To minimize any possible risk, men wishing to father a child should consider discontinuing use of ARAVA and taking cholestyramine 8 grams 3 times daily for 11 days.

Pediatric Use

The safety and effectiveness of ARAVA in pediatric patients with polyarticular course juvenile rheumatoid arthritis (JRA) have not been fully evaluated. (See **CLINICAL STUDIES** and **ADVERSE REACTIONS**).

Geriatric Use

Of the total number of subjects in controlled clinical (Phase III) studies of ARAVA, 234 subjects were 65 years and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out. No dosage adjustment is needed in patients over 65.

ADVERSE REACTIONS

Adverse reactions associated with the use of leflunomide in RA include diarrhea, elevated liver enzymes (ALT and AST), alopecia and rash. In the controlled studies at one year, the following adverse events were reported, regardless of causality. (See Table 9.)

Table 9. Percentage Of Patients With Adverse Events ≥3% In Any Leflunomide Treated Group							
	All RA Studies	Placebo-Controlled Trials				Active-Controlled Trials	
		MN 301 and US 301				MN 302*	
	LEF (N=1339)¹	LEF (N=315)	PBO (N=210)	SSZ (N=133)	MTX (N=182)	LEF (N=501)	MTX (N=498)
BODY AS A WHOLE							
Allergic Reaction	2%	5%	2%	0%	6%	1%	2%
Asthenia	3%	6%	4%	5%	6%	3%	3%
Flu Syndrome	2%	4%	2%	0%	7%	0%	0%
Infection, upper respiratory	4%	0%	0%	0%	0%	0%	0%
Injury Accident	5%	7%	5%	3%	11%	6%	7%
Pain	2%	4%	2%	2%	5%	1%	<1%
Abdominal Pain	6%	5%	4%	4%	8%	6%	4%
Back Pain	5%	6%	3%	4%	9%	8%	7%
CARDIOVASCULAR							
Hypertension ²	10%	9%	4%	4%	3%	10%	4%
- New onset of hypertension		1%	<1%	0%	2%	2%	<1%
Chest Pain	2%	4%	2%	2%	4%	1%	2%
GASTROINTESTINAL							
Anorexia	3%	3%	2%	5%	2%	3%	3%
Diarrhea	17%	27%	12%	10%	20%	22%	10%
Dyspepsia	5%	10%	10%	9%	13%	6%	7%
Gastroenteritis	3%	1%	1%	0%	6%	3%	3%
Abnormal Liver Enzymes	5%	10%	2%	4%	10%	6%	17%
Nausea	9%	13%	11%	19%	18%	13%	18%
GI/Abdominal Pain	5%	6%	4%	7%	8%	8%	8%
Mouth Ulcer	3%	5%	4%	3%	10%	3%	6%
Vomiting	3%	5%	4%	4%	3%	3%	3%
METABOLIC AND NUTRITIONAL							
Hypokalemia	1%	3%	1%	1%	1%	1%	<1%
Weight Loss ³	4%	2%	1%	2%	0%	2%	2%

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

MUSCULO-SKELETAL SYSTEM							
Arthralgia	1%	4%	3%	0%	9%	<1%	1%
Leg Cramps	1%	4%	2%	2%	6%	0%	0%
Joint Disorder	4%	2%	2%	2%	2%	8%	6%
Synovitis	2%	<1%	1%	0%	2%	4%	2%
Tenosynovitis	3%	2%	0%	1%	2%	5%	1%
NERVOUS SYSTEM							
Dizziness	4%	5%	3%	6%	5%	7%	6%
Headache	7%	13%	11%	12%	21%	10%	8%
Paresthesia	2%	3%	1%	1%	2%	4%	3%
RESPIRATORY SYSTEM							
Bronchitis	7%	5%	2%	4%	7%	8%	7%
Increased Cough	3%	4%	5%	3%	6%	5%	7%
Respiratory Infection	15%	21%	21%	20%	32%	27%	25%
Pharyngitis	3%	2%	1%	2%	1%	3%	3%
Pneumonia	2%	3%	0%	0%	1%	2%	2%
Rhinitis	2%	5%	2%	4%	3%	2%	2%
Sinusitis	2%	5%	5%	0%	10%	1%	1%
SKIN AND APPENDAGES							
Alopecia	10%	9%	1%	6%	6%	17%	10%
Eczema	2%	1%	1%	1%	1%	3%	2%
Pruritus	4%	5%	2%	3%	2%	6%	2%
Rash	10%	12%	7%	11%	9%	11%	10%
Dry Skin	2%	3%	2%	2%	0%	3%	1%
UROGENITAL SYSTEM							
Urinary Tract Infection	5%	5%	7%	4%	2%	5%	6%

* Only 10% of patients in MN302 received folate. All patients in US301 received folate; none in MN301 received folate.

- 1 Includes all controlled and uncontrolled trials with leflunomide (duration up to 12 months).
- 2 Hypertension as a preexisting condition was overrepresented in all leflunomide treatment groups in phase III trials
- 3 In a meta-analysis of all phase II and III studies, during the first 6 months in patients receiving leflunomide, 10% lost 10-19 lbs (24 cases per 100 patient years) and 2% lost at least 20 lbs (4 cases/100 patient years). Of patients receiving leflunomide, 4% lost 10% of their baseline weight during the first 6 months of treatment.

Adverse events during a second year of treatment with leflunomide in clinical trials were consistent with those observed during the first year of treatment and occurred at a similar or lower incidence.

In addition, the following adverse events have been reported in 1% to <3% of the RA patients in the leflunomide treatment group in controlled clinical trials.

Body as a Whole: abscess, cyst, fever, hernia, malaise, pain, neck pain, pelvic pain;

Cardiovascular: angina pectoris, migraine, palpitation, tachycardia, varicose vein, vasculitis, vasodilatation;

Gastrointestinal: cholelithiasis, colitis, constipation, esophagitis, flatulence, gastritis, gingivitis, melena, oral moniliasis, pharyngitis, salivary gland enlarged, stomatitis (or aphthous stomatitis), tooth disorder;

Endocrine: diabetes mellitus, hyperthyroidism;

Hemic and Lymphatic System: anemia (including iron deficiency anemia), ecchymosis;

Metabolic and Nutritional: creatine phosphokinase increased, hyperglycemia, hyperlipidemia, peripheral edema;

Musculo-Skeletal System: arthrosis, bone necrosis, bone pain, bursitis, muscle cramps, myalgia, tendon rupture;

Nervous System: anxiety, depression, dry mouth, insomnia, neuralgia, neuritis, sleep disorder, sweating increased, vertigo;

Respiratory System: asthma, dyspnea, epistaxis, lung disorder;

Skin and Appendages: acne, contact dermatitis, fungal dermatitis, hair discoloration, hematoma, herpes simplex, herpes zoster, maculopapular rash, nail disorder, skin discoloration, skin disorder, skin nodule, subcutaneous nodule, ulcer skin;

Special Senses: blurred vision, cataract, conjunctivitis, eye disorder, taste perversion;

Urogenital System: albuminuria, cystitis, dysuria, hematuria, menstrual disorder, prostate disorder, urinary frequency, vaginal moniliasis.

Other less common adverse events seen in clinical trials include: 1 case of anaphylactic reaction occurred in Phase 2 following rechallenge of drug after withdrawal due to rash (rare); urticaria; eosinophilia; transient thrombocytopenia (rare); and leukopenia <2000 WBC/mm³ (rare).

Adverse events during a second year of treatment with leflunomide in clinical trials were consistent with those observed during the first year of treatment and occurred at a similar or lower incidence.

In post-marketing experience, the following have been reported rarely:

Body as a whole: opportunistic infections, severe infections including sepsis that may be fatal;

Gastrointestinal: pancreatitis;

Hematologic: agranulocytosis, leukopenia, neutropenia, pancytopenia, thrombocytopenia;

Hypersensitivity: angioedema;

ARAVA® Tablets
(leflunomide)
10 mg, 20 mg, 100 mg

Hepatic: hepatitis, jaundice/cholestasis, severe liver injury such as hepatic failure and acute hepatic necrosis that may be fatal;

Respiratory: interstitial lung disease, including interstitial pneumonitis and pulmonary fibrosis, which may be fatal;

Nervous system: peripheral neuropathy;

Skin and Appendages: erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis.

Adverse Reactions (Pediatric Patients)

The safety of ARAVA was studied in 74 patients with polyarticular course juvenile rheumatoid arthritis ranging in age from 3-17 years (47 patients from the active-controlled study and 27 from the open-label safety and pharmacokinetic study). The most common adverse events included abdominal pain, diarrhea, nausea, vomiting, oral ulcers, upper respiratory tract infections, alopecia, rash, headache, and dizziness. Less common adverse events included anemia, hypertension, and weight loss. Fourteen pediatric patients experienced ALT and/or AST elevations, nine between 1.2 and 3-fold the upper limit of normal, five between 3 and 8-fold the upper limit of normal.

DRUG ABUSE AND DEPENDENCE

ARAVA has no known potential for abuse or dependence.

OVERDOSAGE

In mouse and rat acute toxicology studies, the minimally toxic dose for oral leflunomide was 200 - 500 mg/kg and 100 mg/kg, respectively (approximately >350 times the maximum recommended human dose, respectively).

There have been reports of chronic overdose in patients taking ARAVA at daily dose up to five times the recommended daily dose and reports of acute overdose in adults or children. There were no adverse events reported in the majority of case reports of overdose. Adverse events were consistent with the safety profile for ARAVA (see **ADVERSE REACTIONS**). The most frequent adverse events observed were diarrhea, abdominal pain, leukopenia, anemia and elevated liver function tests.

In the event of a significant overdose or toxicity, cholestyramine or charcoal administration is recommended to accelerate elimination (see **PRECAUTIONS – General – Need for Drug Elimination**).

Studies with both hemodialysis and CAPD (chronic ambulatory peritoneal dialysis) indicate that M1, the primary metabolite of leflunomide, is not dialyzable. (See **CLINICAL PHARMACOLOGY – Elimination**).

DOSAGE AND ADMINISTRATION

Loading Dose

Due to the long half-life in patients with RA and recommended dosing interval (24 hours), a loading dose is needed to provide steady-state concentrations more rapidly. It is recommended that ARAVA therapy be initiated with a loading dose of one 100 mg tablet per day for 3 days.

Elimination of the loading dose regimen may decrease the risk of adverse events. This could be especially important for patients at increased risk of hematologic or hepatic toxicity, such as those receiving concomitant treatment with methotrexate or other immunosuppressive agents or on such medications in the recent past. (See **WARNINGS – Hepatotoxicity**).

Maintenance Therapy

Daily dosing of 20 mg is recommended for treatment of patients with RA. A small cohort of patients (n=104), treated with 25 mg/day, experienced a greater incidence of side effects; alopecia, weight loss, liver enzyme elevations. Doses higher than 20 mg/day are not recommended. If dosing at 20 mg/day is not well tolerated clinically, the dose may be decreased to 10 mg daily. Liver enzymes must be monitored and dose adjustments may be necessary (see **WARNINGS – Hepatotoxicity**). Due to the prolonged half-life of the active metabolite of leflunomide, patients should be carefully observed after dose reduction, since it may take several weeks for metabolite levels to decline.

HOW SUPPLIED

ARAVA Tablets in 10 and 20 mg strengths are packaged in bottles. ARAVA Tablets 100 mg strength are packaged in blister packs.

ARAVA® (leflunomide) Tablets

Strength	Quantity	NDC Number	Description
10 mg	30 count bottle	0088-2160-30	White, round film-coated tablet embossed with "ZBN" on one side.
20 mg	30 count bottle	0088-2161-30	Light yellow, triangular film-coated tablet embossed with "ZBO" on one side.
100 mg	3 count blister pack	0088-2162-03	White, round film-coated tablet embossed with "ZBP" on one side.

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]. Protect from light.

Rx only.

Rev. November 2004

Manufactured by

Aventis Pharma Specialites, 60200 Compiègne, France
 for

Aventis Pharmaceuticals Inc.

Kansas City, MO 64137

Made in France

©2004 Aventis Pharmaceuticals Inc.

ARA-NOV04-F-A